PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

C07H 21/00 A1	(43) International Publication Date: 17 June 1999 (17.06.99)
(21) International Application Number: PCT/US98/2547 (22) International Filing Date: 1 December 1998 (01.12.9)	(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT,
(30) Priority Data: 08/986,065 5 December 1997 (05.12.97)	Published S With international search report.
 (71) Applicant: NANOGEN, INC. [US/US]; 10398 Pacific Centre Court, San Diego, CA 92121 (US). (72) Inventors: SOSNOWSKI, Ronald, G.; 1013 Adella Avenu Coronado, CA 92118 (US). BUTLER, William, F.; 757 Caloma Circle, Carlsbad, CA 92009 (US). TU, Eugen 3527 Lark Street, San Diego, CA 92103 (US). NEREN BERG,: Michael, I.; 11256 Caminito Inocenta, San Diego CA 92014 (US). HELLER, Michael, J.; 1614 Hawk Vie Drive, Encinitas, CA 92024 (US). EDMAN, Carl, F. 12545—C El Camino Real, San Diego, CA 92130 (US). (74) Agents: MURPHY, David, B. et al.; Lyon & Lyon, LLP, Sui 4700, 633 West Fifth Street, Los Angeles, CA 90071–206 (US). 	e

(54) Title: SELF-ADDRESSABLE SELF-ASSEMBLING MICROELECTRONIC INTEGRATED SYSTEMS, COMPONENT DE-VICES, MECHANISMS, METHODS, AND PROCEDURES FOR MOLECULAR BIOLOGICAL ANALYSIS AND DIAG-NOSTICS

(57) Abstract

A self-adressable, self-assembling microelectronic device is designed and fabricated to actively carry out and control multi-step and multiplex molecular biological reactions in microscopic formats. These reactions include nucleic acid hybridizations, antibody/antigen reactions, diagnostics, and biopolymer synthesis. The device can be fabricated using both micro-lithographic and micro-machining techniques. The device can electronically control the transport and attachment of specific binding entities to specific micro-locations. The specific binding entities include molecular biological molecules such as nucleic acids and polypeptides. The device can subsequently control the transport and reaction of analytes or reactants at the addressed specific micro-locations. The device is able to concentrate analytes and reactants, remove non-specifically bound molecules, provide stringency control for DNA hybridization reactions, and improve the detection of analytes. The device can be electronically replicated.